

Non-linear pulsations in Wolf-Rayet stars

Sebastian Wende, Wolfgang Glatzel, Sonja Schuh

*Institut für Astrophysik, Georg-August-Universität Göttingen,
 Friedrich-Hund-Platz 1, D-37077 Göttingen, Germany*

Abstract. Numerical simulations of the evolution of strange-mode instabilities into the non-linear regime have been performed for a wide range of stellar parameters for Wolf-Rayet stars. It has been shown that the Wolf-Rayet models reach radial velocities which amount to up to 30% of their escape velocity. The acoustic luminosities suggest a connection to the observed mass loss. Most of the models show a *jump* in the mean effective temperature after reaching the non-linear regime. This *jump* is related to the run of the opacity.

To determine the fate of unstable stars, the evolution of their instabilities has to be followed into the non-linear regime by numerical simulations (Grott et al. 2005). Linear stability analyses have shown that strong instabilities with dynamical growth rates prevail in many types of massive stars. These are associated with so called *strange modes* (Gautschy & Glatzel 1990). Models of Wolf-Rayet stars exhibit the strongest *strange mode* instabilities found so far (Glatzel & Kiriakidis 1993). Thus they are promising candidates for a non-linear stability analysis.

The Wolf-Rayet models investigated reach final radial velocity amplitudes in the non-linear regime of the order of 10^7 cm/s . These values are comparable with the escape velocity. To investigate a possible connection between stellar instabilities and a pulsationally driven mass loss, the acoustic energy flux is considered. It represents the mechanical energy transferred by shock waves to the star's atmosphere (Grott et al. 2005). We emphasize that the acoustic luminosity and thus any mass loss rate based on it strongly depends on the artificial viscosity which has to be used in the simulations. However, since viscosity has a dissipative effect the derived mass loss rates have to be regarded as lower limits (see Table 1).

Table 1. Acoustic luminosities and mass loss rates based on them for models with different initial effective temperatures. For the models, the mass-luminosity relation of the He-ZAMS was adopted.

M [$M_{\odot}$]	T_{initial} [K]	$L_{\text{acou.}}$ [erg/s]	$\dot{M}$ [$M_{\odot}/a$]	M [$M_{\odot}$]	T_{initial} [K]	$L_{\text{acou.}}$ [erg/s]	$\dot{M}$ [$M_{\odot}/a$]
7.021	60000	$5.00 \cdot 10^{33}$	$1.40 \cdot 10^{-8}$	13.040	60000	$8.17 \cdot 10^{34}$	$2.30 \cdot 10^{-7}$
7.021	90000	$2.55 \cdot 10^{32}$	$3.17 \cdot 10^{-10}$	13.040	90000	$6.78 \cdot 10^{33}$	$8.44 \cdot 10^{-9}$
9.027	60000	$2.64 \cdot 10^{34}$	$7.50 \cdot 10^{-8}$	17.050	60000	$6.84 \cdot 10^{34}$	$1.87 \cdot 10^{-7}$
9.027	90000	$1.54 \cdot 10^{33}$	$1.95 \cdot 10^{-9}$	17.050	90000	$1.95 \cdot 10^{35}$	$2.37 \cdot 10^{-7}$

After reaching the non-linear regime all unstable models exhibit a *jump* in the mean effective temperature. The final mean effective temperature is independent of mass and largely independent of the initial effective temperature. We find only two final mean effective temperatures around $30000K$ and $15000K$ respectively. Models with an initial effective temperature above $\approx 32000K$ reach a mean effective temperature around $30000K$, models with initial temperatures below this value end up with a mean effective temperature around $15000K$. Considering the run of the opacity, these two temperatures correspond to the descending branches of the opacity from the respective maximum towards lower temperatures (see Fig.1). Should descending branches of the opacity in general give rise to a range of possible final mean temperatures of pulsating models, a third range above $\approx 100000K$ would be expected. In fact, some models show a *jump* to this temperature. The final mean position of the pulsating models in the HRD is of particular interest. Most of the models investigated move to the right and settle in a narrow temperature strip (Fig.1). It is remarkably close to the observed positions of Wolf-Rayet stars.




Figure 1. Left: Run of the opacity in the stellar envelope as a function of temperature. Different stellar models were used (different gray scales correspond to different models) to cover a wide range of temperatures. For further details see text. Right: Hertzsprung-Russell-Diagram. Indicated are the positions of the initial models by crosses and asterisks and the final mean positions by squares and circles. The mean luminosities stay constant. Arrows indicate the direction of the evolution. For models with question marks, we were not able to determine a final mean temperature. Models without arrows don't change their mean temperature. The He-ZAMS is shown for comparison.

Acknowledgments. Financial support by the Institut für Astronomie und Astrophysik Tübingen is gratefully acknowledged.

References

- Grott, M., Chernigovski, S. & Glatzel, W. 2005, MNRAS, 360, 1532
 Gautschi, A. & Glatzel, W. 1990, MNRAS, 245, 597
 Glatzel, W. & Kiriakidis, M. 1993, MNRAS, 262, L7